

The geometry of the cyclotomic trace

Aaron Mazel-Gee

with David Ayala and Nick Rozenblyum

X a scheme (derived)

X a scheme (derived)

$K(X) =$ *algebraic K-theory* of X

X a scheme (derived)

$K(X) =$ *algebraic K-theory* of X
 $:= K(\text{Perf}_X)$

X a scheme (derived)

$K(X) =$ *algebraic K-theory* of X

$:= K(\text{Perf}_X)$

\approx group-completion of $(\text{VBdl}(X)/\text{iso.}, \oplus)$

X a scheme (derived)

$K(X) =$ *algebraic K-theory* of X

$:= K(\text{Perf}_X)$

\approx group-completion of $(\text{VBdl}(X)/\text{iso.}, \oplus)$

[hard to compute!]

X a scheme (derived)

$K(X) =$ *algebraic K-theory* of X

$:= K(\text{Perf}_X)$

\approx group-completion of $(\text{VBdl}(X)/\text{iso.}, \oplus)$

[hard to compute!]

$\text{THH}(X) =$ *topological Hochschild homology* of X

X a scheme (derived)

$K(X) =$ *algebraic K-theory* of X

$:= K(\text{Perf}_X)$

\approx group-completion of $(\text{VBdl}(X)/\text{iso.}, \oplus)$

[hard to compute!]

$\text{THH}(X) =$ *topological Hochschild homology* of X

$:= \text{THH}(\text{Perf}_X)$

X a scheme (derived)

$K(X) =$ *algebraic K-theory* of X

$:= K(\text{Perf}_X)$

\approx group-completion of $(\text{VBdl}(X)/\text{iso.}, \oplus)$

[hard to compute!]

$\text{THH}(X) =$ *topological Hochschild homology* of X

$:= \text{THH}(\text{Perf}_X) := \int_{S^1} \text{Perf}_X$

X a scheme (derived)

$K(X) =$ *algebraic K-theory* of X

$:= K(\text{Perf}_X)$

\approx group-completion of $(\text{VBdl}(X)/\text{iso.}, \oplus)$

[hard to compute!]

$\text{THH}(X) =$ *topological Hochschild homology* of X

$:= \text{THH}(\text{Perf}_X) := \int_{S^1} \text{Perf}_X$

$\approx \mathcal{O}(\mathcal{L}X)$, functions on the free loop space of X

X a scheme (derived)

$K(X) =$ *algebraic K-theory* of X

$:= K(\text{Perf}_X)$

\approx group-completion of $(\text{VBdl}(X)/\text{iso.}, \oplus)$

[hard to compute!]

$\text{THH}(X) =$ *topological Hochschild homology* of X

$:= \text{THH}(\text{Perf}_X) := \int_{S^1} \text{Perf}_X$

$\approx \mathcal{O}(\mathcal{L}X)$, functions on the free loop space of X

[easier to compute]

X a scheme (derived)

$K(X) =$ *algebraic K-theory* of X

$$:= K(\text{Perf}_X)$$

\approx group-completion of $(\text{VBdl}(X)/\text{iso.}, \oplus)$

[hard to compute!]

$\text{THH}(X) =$ *topological Hochschild homology* of X

$$:= \text{THH}(\text{Perf}_X) := \int_{S^1} \text{Perf}_X$$

$\approx \mathcal{O}(\mathcal{L}X)$, functions on the free loop space of X

[easier to compute]

the *Dennis trace* map

$$K(X) \longrightarrow \text{THH}(X)$$

X a scheme (derived)

$K(X) =$ *algebraic K-theory* of X

$$:= K(\text{Perf}_X)$$

$$\approx \text{group-completion of } (\text{VBdl}(X)/\text{iso.}, \oplus)$$

[hard to compute!]

$\text{THH}(X) =$ *topological Hochschild homology* of X

$$:= \text{THH}(\text{Perf}_X) := \int_{S^1} \text{Perf}_X$$

$$\approx \mathcal{O}(\mathcal{L}X), \text{ functions on the free loop space of } X$$

[easier to compute]

the *Dennis trace* map

$$K(X) \longrightarrow \text{THH}(X)$$

$$(E \downarrow X) \longmapsto \left(\begin{array}{c} \\ \\ \\ \end{array} \right)$$

X a scheme (derived)

$K(X) =$ *algebraic K-theory* of X

$$:= K(\text{Perf}_X)$$

$$\approx \text{group-completion of } (\text{VBdl}(X)/\text{iso.}, \oplus)$$

[hard to compute!]

$\text{THH}(X) =$ *topological Hochschild homology* of X

$$:= \text{THH}(\text{Perf}_X) := \int_{S^1} \text{Perf}_X$$

$$\approx \mathcal{O}(\mathcal{L}X), \text{ functions on the free loop space of } X$$

[easier to compute]

the *Dennis trace* map

$$K(X) \longrightarrow \text{THH}(X)$$

$$(E \downarrow X) \longmapsto \left((S^1 \xrightarrow{\gamma} X) \longmapsto \right)$$

X a scheme (derived)

$K(X) =$ *algebraic K-theory* of X

$$:= K(\text{Perf}_X)$$

$$\approx \text{group-completion of } (\text{VBdl}(X)/\text{iso.}, \oplus)$$

[hard to compute!]

$\text{THH}(X) =$ *topological Hochschild homology* of X

$$:= \text{THH}(\text{Perf}_X) := \int_{S^1} \text{Perf}_X$$

$$\approx \mathcal{O}(\mathcal{L}X), \text{ functions on the free loop space of } X$$

[easier to compute]

the *Dennis trace* map

$$K(X) \longrightarrow \text{THH}(X)$$

$$(E \downarrow X) \longmapsto \left((S^1 \xrightarrow{\gamma} X) \longmapsto \begin{array}{l} \text{trace of} \\ \text{monodromy} \\ \text{of } E|_{\gamma} \end{array} \right)$$

a refinement:

$$\begin{array}{ccc} K(X) & \xrightarrow{\text{Dennis trace}} & THH(X) \\ & \searrow \text{cyclotomic trace} & \nearrow \\ & TC(X) & \end{array}$$

a refinement:

$$\begin{array}{ccc} K(X) & \xrightarrow{\text{Dennis trace}} & THH(X) \\ & \searrow \text{cyclotomic trace} & \nearrow \\ & TC(X) & \end{array}$$

$TC(X) = \textit{topological cyclic homology}$ of X

a refinement:

$$\begin{array}{ccc} K(X) & \xrightarrow{\text{Dennis trace}} & THH(X) \\ & \searrow \text{cyclotomic trace} & \nearrow \\ & TC(X) & \end{array}$$

$TC(X) = \textit{topological cyclic homology}$ of X
 $\approx \dots???!?!?$

a refinement:

$$\begin{array}{ccc} K(X) & \xrightarrow{\text{Dennis trace}} & THH(X) \\ & \searrow \text{cyclotomic trace} & \nearrow \\ & TC(X) & \end{array}$$

$TC(X) = \textit{topological cyclic homology}$ of X
 $\approx \dots???!?!?$

[computationally accessible, but conceptually mysterious]

why we care about TC

Theorem (Goodwillie/ \mathbb{Q} '86, McCarthy/ \mathbb{Z} '97, Dundas/ \mathbb{S} '97)

why we care about TC

Theorem (Goodwillie/ \mathbb{Q} '86, McCarthy/ \mathbb{Z} '97, Dundas/ \mathbb{S} '97)

The cyclotomic trace is “locally constant”:

why we care about TC

Theorem (Goodwillie/ \mathbb{Q} '86, McCarthy/ \mathbb{Z} '97, Dundas/ \mathbb{S} '97)

The cyclotomic trace is “locally constant”: for $\tilde{A} \rightarrow A$ a nilpotent extension of associative rings (or of connective ring spectra),

why we care about TC

Theorem (Goodwillie/ \mathbb{Q} '86, McCarthy/ \mathbb{Z} '97, Dundas/ \mathbb{S} '97)

The cyclotomic trace is “locally constant”: for $\tilde{A} \rightarrow A$ a nilpotent extension of associative rings (or of connective ring spectra),

$$\begin{array}{ccc} K(\tilde{A}) & \longrightarrow & K(A) \\ \downarrow & & \downarrow \\ TC(\tilde{A}) & \longrightarrow & TC(A) \end{array}$$

is a pullback (after p -completion, for any prime p).

why we care about TC

Theorem (Goodwillie/ \mathbb{Q} '86, McCarthy/ \mathbb{Z} '97, Dundas/ \mathbb{S} '97)

The cyclotomic trace is “locally constant”: for $\tilde{A} \rightarrow A$ a nilpotent extension of associative rings (or of connective ring spectra),

$$\begin{array}{ccc} K(\tilde{A}) & \longrightarrow & K(A) \\ \downarrow & & \downarrow \\ TC(\tilde{A}) & \longrightarrow & TC(A) \end{array}$$

is a pullback (after p -completion, for any prime p).

“This is how people other than Quillen compute algebraic K-theory.”

~ A. Blumberg, algebraic K-theorist

...but what *is* $TC(X)$, really?

...but what *is* $TC(X)$, really?

intermediate factorization through *negative cyclic homology*:

$$K(X) \longrightarrow TC(X) \longrightarrow HC^-(X) \longrightarrow THH(X)$$

...but what *is* $TC(X)$, really?

intermediate factorization through *negative cyclic homology*:

$$K(X) \longrightarrow TC(X) \longrightarrow HC^-(X) \longrightarrow THH(X)$$

	differential algebra	derived algebraic geometry
$THH(X)$	$\Omega_{dR}^*(X)$	functions on $\mathcal{L}X$

...but what *is* $TC(X)$, really?

intermediate factorization through *negative cyclic homology*:

$$K(X) \longrightarrow TC(X) \longrightarrow HC^-(X) \longrightarrow THH(X)$$

	differential algebra	derived algebraic geometry
$THH(X)$	$\Omega_{dR}^*(X)$	functions on $\mathcal{L}X$
$HC^-(X)$ $:= THH(X)^{h\mathbb{T}}$		

...but what *is* $TC(X)$, really?

intermediate factorization through *negative cyclic homology*:

$$K(X) \longrightarrow TC(X) \longrightarrow HC^-(X) \longrightarrow THH(X)$$

	differential algebra	derived algebraic geometry
$THH(X)$	$\Omega_{dR}^*(X)$	functions on $\mathcal{L}X$
$HC^-(X)$ $:= THH(X)^{h\mathbb{T}}$	$H_{dR}^*(X)$	

...but what *is* $TC(X)$, really?

intermediate factorization through *negative cyclic homology*:

$$K(X) \longrightarrow TC(X) \longrightarrow HC^-(X) \longrightarrow THH(X)$$

	differential algebra	derived algebraic geometry
$THH(X)$	$\Omega_{dR}^*(X)$	functions on $\mathcal{L}X$
$HC^-(X)$ $:= THH(X)^{h\mathbb{T}}$	$H_{dR}^*(X)$	\mathbb{T} -invariant functions on $\mathcal{L}X$

...but what *is* $TC(X)$, really?

intermediate factorization through *negative cyclic homology*:

$$K(X) \longrightarrow TC(X) \longrightarrow HC^-(X) \longrightarrow THH(X)$$

	differential algebra	derived algebraic geometry
$THH(X)$	$\Omega_{dR}^*(X)$	functions on $\mathcal{L}X$
$HC^-(X)$ $:= THH(X)^{h\mathbb{T}}$	$H_{dR}^*(X)$	\mathbb{T} -invariant functions on $\mathcal{L}X$
$TC(X)$ $:= THH(X)^{hCyc}$		

...but what *is* $TC(X)$, really?

intermediate factorization through *negative cyclic homology*:

$$K(X) \longrightarrow TC(X) \longrightarrow HC^-(X) \longrightarrow THH(X)$$

	differential algebra	derived algebraic geometry
$THH(X)$	$\Omega_{dR}^*(X)$	functions on $\mathcal{L}X$
$HC^-(X)$ $:= THH(X)^{h\mathbb{T}}$	$H_{dR}^*(X)$	\mathbb{T} -invariant functions on $\mathcal{L}X$
$TC(X)$ $:= THH(X)^{hCyc}$	$???_{dR}^*(X)$	

...but what *is* $TC(X)$, really?

intermediate factorization through *negative cyclic homology*:

$$K(X) \longrightarrow TC(X) \longrightarrow HC^-(X) \longrightarrow THH(X)$$

	differential algebra	derived algebraic geometry
$THH(X)$	$\Omega_{dR}^*(X)$	functions on $\mathcal{L}X$
$HC^-(X)$ $:= THH(X)^{h\mathbb{T}}$	$H_{dR}^*(X)$	\mathbb{T} -invariant functions on $\mathcal{L}X$
$TC(X)$ $:= THH(X)^{hCyc}$	$???_{dR}^*(X)$	TODAY

constructions of TC

constructions of TC

original definition (Bökstedt–Hsiang–Madsen '93):

constructions of TC

original definition (Bökstedt–Hsiang–Madsen '93):

- uses *genuine-equivariant* stable homotopy theory

constructions of TC

original definition (Bökstedt–Hsiang–Madsen '93):

- uses *genuine-equivariant* stable homotopy theory
 - useful (e.g. equivariant Poincaré duality)...

constructions of TC

original definition (Bökstedt–Hsiang–Madsen '93):

- uses *genuine-equivariant* stable homotopy theory
 - useful (e.g. equivariant Poincaré duality)...
 - but not conceptual (no DAG interpretation known)

constructions of TC

original definition (Bökstedt–Hsiang–Madsen '93):

- uses *genuine-equivariant* stable homotopy theory
 - useful (e.g. equivariant Poincaré duality)...
 - but not conceptual (no DAG interpretation known)
- used opaque point-set manipulations

constructions of TC

original definition (Bökstedt–Hsiang–Madsen '93):

- uses *genuine-equivariant* stable homotopy theory
 - useful (e.g. equivariant Poincaré duality)...
 - but not conceptual (no DAG interpretation known)
- used opaque point-set manipulations
- based on vague analogy with free loopspaces

constructions of TC

original definition (Bökstedt–Hsiang–Madsen '93):

- uses *genuine-equivariant* stable homotopy theory
 - useful (e.g. equivariant Poincaré duality)...
 - but not conceptual (no DAG interpretation known)
- used opaque point-set manipulations
- based on vague analogy with free loopspaces

firmer categorical footing (Blumberg–Mandell '13):

- define homotopy theory of “cyclotomic spectra”

constructions of TC

original definition (Bökstedt–Hsiang–Madsen '93):

- uses *genuine-equivariant* stable homotopy theory
 - useful (e.g. equivariant Poincaré duality)...
 - but not conceptual (no DAG interpretation known)
- used opaque point-set manipulations
- based on vague analogy with free loopspaces

firmer categorical footing (Blumberg–Mandell '13):

- define homotopy theory of “cyclotomic spectra”

more recent definition (Nikolaus–Scholze '17?):

constructions of TC

original definition (Bökstedt–Hsiang–Madsen '93):

- uses *genuine-equivariant* stable homotopy theory
 - useful (e.g. equivariant Poincaré duality)...
 - but not conceptual (no DAG interpretation known)
- used opaque point-set manipulations
- based on vague analogy with free loopspaces

firmer categorical footing (Blumberg–Mandell '13):

- define homotopy theory of “cyclotomic spectra”

more recent definition (Nikolaus–Scholze '17?):

- removes genuine-equivariance

constructions of TC

original definition (Bökstedt–Hsiang–Madsen '93):

- uses *genuine-equivariant* stable homotopy theory
 - useful (e.g. equivariant Poincaré duality)...
 - but not conceptual (no DAG interpretation known)
- used opaque point-set manipulations
- based on vague analogy with free loopspaces

firmer categorical footing (Blumberg–Mandell '13):

- define homotopy theory of “cyclotomic spectra”

more recent definition (Nikolaus–Scholze '17?):

- removes genuine-equivariance
- restricts to *connective* ring spectra

constructions of TC

original definition (Bökstedt–Hsiang–Madsen '93):

- uses *genuine-equivariant* stable homotopy theory
 - useful (e.g. equivariant Poincaré duality)...
 - but not conceptual (no DAG interpretation known)
- used opaque point-set manipulations
- based on vague analogy with free loopspaces

firmer categorical footing (Blumberg–Mandell '13):

- define homotopy theory of “cyclotomic spectra”

more recent definition (Nikolaus–Scholze '17?):

- removes genuine-equivariance
- restricts to *connective* ring spectra

this talk, inspired by Nikolaus–Scholze:

constructions of TC

original definition (Bökstedt–Hsiang–Madsen '93):

- uses *genuine-equivariant* stable homotopy theory
 - useful (e.g. equivariant Poincaré duality)...
 - but not conceptual (no DAG interpretation known)
- used opaque point-set manipulations
- based on vague analogy with free loopspaces

firmer categorical footing (Blumberg–Mandell '13):

- define homotopy theory of “cyclotomic spectra”

more recent definition (Nikolaus–Scholze '17?):

- removes genuine-equivariance
- restricts to *connective* ring spectra

this talk, inspired by Nikolaus–Scholze:

- applies to any spectrally-enriched ∞ -category

constructions of TC

original definition (Bökstedt–Hsiang–Madsen '93):

- uses *genuine-equivariant* stable homotopy theory
 - useful (e.g. equivariant Poincaré duality)...
 - but not conceptual (no DAG interpretation known)
- used opaque point-set manipulations
- based on vague analogy with free loopspaces

firmer categorical footing (Blumberg–Mandell '13):

- define homotopy theory of “cyclotomic spectra”

more recent definition (Nikolaus–Scholze '17?):

- removes genuine-equivariance
- restricts to *connective* ring spectra

this talk, inspired by Nikolaus–Scholze:

- applies to any spectrally-enriched ∞ -category
- uses *factorization homology* to keep track of symmetries

constructions of TC

original definition (Bökstedt–Hsiang–Madsen '93):

- uses *genuine-equivariant* stable homotopy theory
 - useful (e.g. equivariant Poincaré duality)...
 - but not conceptual (no DAG interpretation known)
- used opaque point-set manipulations
- based on vague analogy with free loopspaces

firmer categorical footing (Blumberg–Mandell '13):

- define homotopy theory of “cyclotomic spectra”

more recent definition (Nikolaus–Scholze '17?):

- removes genuine-equivariance
- restricts to *connective* ring spectra

this talk, inspired by Nikolaus–Scholze:

- applies to any spectrally-enriched ∞ -category
- uses *factorization homology* to keep track of symmetries
- admits direct interpretation in DAG via $\mathcal{L}X$

constructions of TC

original definition (Bökstedt–Hsiang–Madsen '93):

- uses *genuine-equivariant* stable homotopy theory
 - useful (e.g. equivariant Poincaré duality)...
 - but not conceptual (no DAG interpretation known)
- used opaque point-set manipulations
- based on vague analogy with free loopspaces

firmer categorical footing (Blumberg–Mandell '13):

- define homotopy theory of “cyclotomic spectra”

more recent definition (Nikolaus–Scholze '17?):

- removes genuine-equivariance
- restricts to *connective* ring spectra

this talk, inspired by Nikolaus–Scholze:

- applies to any spectrally-enriched ∞ -category
- uses *factorization homology* to keep track of symmetries
- admits direct interpretation in DAG via $\mathcal{L}X$
- suggests higher-dim generalizations (\rightsquigarrow “higher K-theory”)

overview

overview

$$\begin{array}{ccc} \mathrm{Sp} & \xrightarrow{\mathrm{triv}} & \mathrm{Cyc}(\mathrm{Sp}) \\ & \xleftarrow{\perp} & \\ & \xleftarrow{(-)^{\mathrm{hCyc}}} & \\ \Psi & & \Psi \end{array}$$
$$\mathrm{TC}(X) \longleftarrow \mathrm{THH}(X)$$

overview

$$\begin{array}{ccc} \mathrm{Sp} & \xrightarrow{\mathrm{triv}} & \mathrm{Cyc}(\mathrm{Sp}) \\ \downarrow \Psi & \xleftarrow[(-)^{\mathrm{hCyc}}]{\perp} & \downarrow \Psi \\ \mathrm{TC}(X) & \longleftarrow \longrightarrow & \mathrm{THH}(X) \end{array}$$

$\mathrm{TC}(X)$:= fixedpoints of *cyclotomic structure* on $\mathrm{THH}(X)$
 \rightsquigarrow built by “imposing conditions” on functions on $\mathcal{L}X$

overview

$$\begin{array}{ccc} \mathrm{Sp} & \xrightarrow{\mathrm{triv}} & \mathrm{Cyc}(\mathrm{Sp}) \\ & \xleftarrow{\perp} & \\ & \xleftarrow{(-)^{\mathrm{hCyc}}} & \\ \Psi & & \Psi \end{array}$$
$$\mathrm{TC}(X) \longleftarrow \mathrm{THH}(X)$$

$\mathrm{TC}(X)$:= fixedpoints of *cyclotomic structure* on $\mathrm{THH}(X)$
 \rightsquigarrow built by “imposing conditions” on functions on $\mathcal{L}X$

main idea: $\mathrm{TC}(X) \approx$ functions on $\mathcal{L}X$ that are...

overview

$$\begin{array}{ccc} \mathrm{Sp} & \xrightarrow{\mathrm{triv}} & \mathrm{Cyc}(\mathrm{Sp}) \\ & \xleftarrow{\perp} & \\ & \xleftarrow{(-)^{\mathrm{hCyc}}} & \\ \Psi & & \Psi \end{array}$$
$$\mathrm{TC}(X) \longleftarrow \mathrm{THH}(X)$$

$\mathrm{TC}(X)$:= fixedpoints of *cyclotomic structure* on $\mathrm{THH}(X)$
 \rightsquigarrow built by “imposing conditions” on functions on $\mathcal{L}X$

main idea: $\mathrm{TC}(X) \approx$ functions on $\mathcal{L}X$ that are...

- invariant under the \mathbb{T} -action on $\mathcal{L}X$.

overview

$$\begin{array}{ccc} \mathrm{Sp} & \xrightarrow{\mathrm{triv}} & \mathrm{Cyc}(\mathrm{Sp}) \\ & \xleftarrow{\perp} & \\ & \xleftarrow{(-)^{\mathrm{hCyc}}} & \\ \Psi & & \Psi \end{array}$$
$$\mathrm{TC}(X) \longleftarrow \mathrm{THH}(X)$$

$\mathrm{TC}(X) :=$ fixedpoints of *cyclotomic structure* on $\mathrm{THH}(X)$
 \rightsquigarrow built by “imposing conditions” on functions on $\mathcal{L}X$

main idea: $\mathrm{TC}(X) \approx$ functions on $\mathcal{L}X$ that are...

- invariant under the \mathbb{T} -action on $\mathcal{L}X$.
- “sensitive” to the relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$.

overview

$$\begin{array}{ccc} \mathrm{Sp} & \xrightarrow{\mathrm{triv}} & \mathrm{Cyc}(\mathrm{Sp}) \\ & \xleftarrow{\perp} & \\ & \xleftarrow{(-)^{\mathrm{hCyc}}} & \\ \Psi & & \Psi \end{array}$$
$$\mathrm{TC}(X) \longleftarrow \mathrm{THH}(X)$$

$\mathrm{TC}(X)$:= fixedpoints of *cyclotomic structure* on $\mathrm{THH}(X)$
 \rightsquigarrow built by “imposing conditions” on functions on $\mathcal{L}X$

main idea: $\mathrm{TC}(X) \approx$ functions on $\mathcal{L}X$ that are...

- invariant under the \mathbb{T} -action on $\mathcal{L}X$.
- “sensitive” to the relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$.

\mathbb{T} -invariance is easy, but what does “sensitive” mean?

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

Q.: M an $n \times n$ matrix, compare $\text{tr}(M)^r$ and $\text{tr}(M^r)$?

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

Q.: M an $n \times n$ matrix, compare $\text{tr}(M)^r$ and $\text{tr}(M^r)$?

Ex. 1: $r = 2$, $M = \text{diag}(m_1, \dots, m_n) \in M_{n \times n}(R)$

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

Q.: M an $n \times n$ matrix, compare $\text{tr}(M)^r$ and $\text{tr}(M^r)$?

Ex. 1: $r = 2$, $M = \text{diag}(m_1, \dots, m_n) \in M_{n \times n}(R)$

$$\text{tr}(M)^2 = \sum_{i,j} m_i m_j \quad ,$$

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

Q.: M an $n \times n$ matrix, compare $\text{tr}(M)^r$ and $\text{tr}(M^r)$?

Ex. 1: $r = 2$, $M = \text{diag}(m_1, \dots, m_n) \in M_{n \times n}(R)$

$$\text{tr}(M)^2 = \sum_{i,j} m_i m_j \quad , \quad \text{tr}(M^2) = \sum_k (m_k)^2$$

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

Q.: M an $n \times n$ matrix, compare $\text{tr}(M)^r$ and $\text{tr}(M^r)$?

Ex. 1: $r = 2$, $M = \text{diag}(m_1, \dots, m_n) \in M_{n \times n}(R)$

$$\text{tr}(M)^2 = \sum_{i,j} m_i m_j \quad , \quad \text{tr}(M^2) = \sum_k (m_k)^2$$

- both *cyclically invariant*, i.e. lie in the fixedpoints $(R \otimes R)^{C_2}$

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

Q.: M an $n \times n$ matrix, compare $\text{tr}(M)^r$ and $\text{tr}(M^r)$?

Ex. 1: $r = 2$, $M = \text{diag}(m_1, \dots, m_n) \in M_{n \times n}(R)$

$$\text{tr}(M)^2 = \sum_{i,j} m_i m_j \quad , \quad \text{tr}(M^2) = \sum_k (m_k)^2$$

- both *cyclically invariant*, i.e. lie in the fixedpoints $(R \otimes R)^{C_2}$
- difference is *norms*:

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

Q.: M an $n \times n$ matrix, compare $\text{tr}(M)^r$ and $\text{tr}(M^r)$?

Ex. 1: $r = 2$, $M = \text{diag}(m_1, \dots, m_n) \in M_{n \times n}(R)$

$$\text{tr}(M)^2 = \sum_{i,j} m_i m_j \quad , \quad \text{tr}(M^2) = \sum_k (m_k)^2$$

- both *cyclically invariant*, i.e. lie in the fixedpoints $(R \otimes R)^{C_2}$
- difference is *norms*: image of $\sum_{i < j} [m_i \otimes m_j]$ under

$$(R \otimes R)_{C_2} \xrightarrow{\text{Nm}} (R \otimes R)^{C_2}$$
$$[x \otimes y] \mapsto \sum_{\sigma \in C_2} \sigma(x \otimes y)$$

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

Q.: M an $n \times n$ matrix, compare $\text{tr}(M)^r$ and $\text{tr}(M^r)$?

Ex. 1: $r = 2$, $M = \text{diag}(m_1, \dots, m_n) \in M_{n \times n}(R)$

$$\text{tr}(M)^2 = \sum_{i,j} m_i m_j \quad , \quad \text{tr}(M^2) = \sum_k (m_k)^2$$

- both *cyclically invariant*, i.e. lie in the fixedpoints $(R \otimes R)^{C_2}$
- difference is *norms*: image of $\sum_{i < j} [m_i \otimes m_j]$ under

$$(R \otimes R)_{C_2} \xrightarrow{\text{Nm}} (R \otimes R)^{C_2}$$
$$[x \otimes y] \mapsto \sum_{\sigma \in C_2} \sigma(x \otimes y)$$

\rightsquigarrow become equal in the **Tate construction**, the cofiber

$$(R \otimes R)_{C_2} \xrightarrow{\text{Nm}} (R \otimes R)^{C_2} \longrightarrow (R \otimes R)^{tC_2}$$

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

Q.: M an $n \times n$ matrix, compare $\text{tr}(M)^r$ and $\text{tr}(M^r)$?

Ex. 2:

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

Q.: M an $n \times n$ matrix, compare $\text{tr}(M)^r$ and $\text{tr}(M^r)$?

Ex. 2: $M = \text{diag}(m_1, m_2)$, $r \in \mathbb{N}^\times$ arbitrary

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

Q.: M an $n \times n$ matrix, compare $\text{tr}(M)^r$ and $\text{tr}(M^r)$?

Ex. 2: $M = \text{diag}(m_1, m_2)$, $r \in \mathbb{N}^\times$ arbitrary

now, difference between

$$\text{tr}(M)^r = (m_1 + m_2)^r, \quad \text{tr}(M^r) = ((m_1)^r + (m_2)^r)$$

governed by binomial coefficients $\binom{r}{i}$ for $0 < i < r$

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

Q.: M an $n \times n$ matrix, compare $\text{tr}(M)^r$ and $\text{tr}(M^r)$?

Ex. 2: $M = \text{diag}(m_1, m_2)$, $r \in \mathbb{N}^\times$ arbitrary

now, difference between

$$\text{tr}(M)^r = (m_1 + m_2)^r, \quad \text{tr}(M^r) = ((m_1)^r + (m_2)^r)$$

governed by binomial coefficients $\binom{r}{i}$ for $0 < i < r$

fact: these are never coprime to r

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

Q.: M an $n \times n$ matrix, compare $\text{tr}(M)^r$ and $\text{tr}(M^r)$?

Ex. 2: $M = \text{diag}(m_1, m_2)$, $r \in \mathbb{N}^\times$ arbitrary

now, difference between

$$\text{tr}(M)^r = (m_1 + m_2)^r, \quad \text{tr}(M^r) = ((m_1)^r + (m_2)^r)$$

governed by binomial coefficients $\binom{r}{i}$ for $0 < i < r$

fact: these are never coprime to r

\rightsquigarrow quotient $(R^{\otimes r})^{C_r}$ by norms from *all* proper subgroups of C_r

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

Q.: M an $n \times n$ matrix, compare $\text{tr}(M)^r$ and $\text{tr}(M^r)$?

Ex. 2: $M = \text{diag}(m_1, m_2)$, $r \in \mathbb{N}^\times$ arbitrary

now, difference between

$$\text{tr}(M)^r = (m_1 + m_2)^r, \quad \text{tr}(M^r) = ((m_1)^r + (m_2)^r)$$

governed by binomial coefficients $\binom{r}{i}$ for $0 < i < r$

fact: these are never coprime to r

\rightsquigarrow quotient $(R^{\otimes r})^{C_r}$ by norms from *all* proper subgroups of C_r

$\rightsquigarrow \text{tr}(M^r) \equiv \text{tr}(M)^r$ in the *generalized* Tate construction $(R^{\otimes r})^{\tau C_r}$

relationship between $S^1 \xrightarrow{\gamma} X$ and $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$

Q.: M an $n \times n$ matrix, compare $\text{tr}(M)^r$ and $\text{tr}(M^r)$?

Ex. 2: $M = \text{diag}(m_1, m_2)$, $r \in \mathbb{N}^\times$ arbitrary

now, difference between

$\text{tr}(M)^r = (m_1 + m_2)^r$, $\text{tr}(M^r) = ((m_1)^r + (m_2)^r)$
governed by binomial coefficients $\binom{r}{i}$ for $0 < i < r$

fact: these are never coprime to r

\rightsquigarrow quotient $(R^{\otimes r})^{C_r}$ by norms from *all* proper subgroups of C_r

$\rightsquigarrow \text{tr}(M^r) \equiv \text{tr}(M)^r$ in the *generalized* Tate construction $(R^{\otimes r})^{\tau C_r}$

★ for \mathcal{C} a spectrally enriched ∞ -category, a covering map

$$S_b^1 \xleftarrow{r} S_a^1$$

of oriented circles induces a ***cyclotomic structure map***

$$\text{THH}(\mathcal{C}) := \int_{S_b^1} \mathcal{C} \longrightarrow \left(\int_{S_a^1} \mathcal{C} \right)^{\tau C_r} =: \text{THH}(\mathcal{C})^{\tau C_r}$$

Theorem 1 (A & M-G & R)

Theorem 1 (A & M-G & R)

$$\text{Cyc}(\text{Sp}) \simeq$$

Theorem 1 (A & M-G & R)

$$\text{Cyc}(\text{Sp}) \simeq \left(\text{Fun}(B\mathbb{T}, \text{Sp}) \right)$$

Theorem 1 (A & M-G & R)

$$\text{Cyc}(\text{Sp}) \simeq \left(\text{Fun}(\text{BT}, \text{Sp}) \underset{\tau}{\curvearrowright} \mathbb{N}^\times \right)$$

Theorem 1 (A & M-G & R)

$$\text{Cyc}(\text{Sp}) \simeq \left(\text{Fun}(\text{BT}, \text{Sp}) \underset{\tau}{\overset{\text{l.lax}}{\curvearrowright}} \mathbb{N}^\times \right)$$

Theorem 1 (A & M-G & R)

$$\text{Cyc}(\text{Sp}) \simeq \lim \left(\text{Fun}(\text{BT}, \text{Sp}) \begin{array}{c} \xrightarrow{\text{lax}} \\ \tau \\ \xleftarrow{\quad} \end{array} \mathbb{N}^\times \right)$$

Theorem 1 (A & M-G & R)

$$\text{Cyc}(\text{Sp}) \simeq \lim^{r.\text{lax}} \left(\text{Fun}(\text{BT}, \text{Sp}) \begin{array}{c} \xrightarrow{\text{l.lax}} \\ \tau \\ \xleftarrow{\text{l.lax}} \end{array} \mathbb{N}^\times \right)$$

Theorem 1 (A & M-G & R)

$$\text{Cyc}(\text{Sp}) \simeq \lim^{r.\text{lax}} \left(\text{Fun}(\text{BT}, \text{Sp}) \underset{\tau}{\overset{l.\text{lax}}{\curvearrowright}} \mathbb{N}^\times \right).$$

Theorem 1 (A & M-G & R)

$$\text{Cyc}(\text{Sp}) \simeq \lim^{r.\text{lax}} \left(\text{Fun}(\mathbb{B}\mathbb{T}, \text{Sp}) \underset{\tau}{\overset{l.\text{lax}}{\curvearrowright}} \mathbb{N}^\times \right).$$

★ an object of $\lim^{r.\text{lax}}$ is given by $T \in \text{Fun}(\mathbb{B}\mathbb{T}, \text{Sp})$ equipped with:

Theorem 1 (A & M-G & R)

$$\mathrm{Cyc}(\mathrm{Sp}) \simeq \lim^{r.\mathrm{lax}} \left(\mathrm{Fun}(\mathrm{BT}, \mathrm{Sp}) \begin{array}{c} \xrightarrow{l.\mathrm{lax}} \\ \tau \\ \xleftarrow{l.\mathrm{lax}} \end{array} \mathbb{N}^\times \right).$$

- ★ an object of $\lim^{r.\mathrm{lax}}$ is given by $T \in \mathrm{Fun}(\mathrm{BT}, \mathrm{Sp})$ equipped with:
- for each $r \in \mathbb{N}^\times$,

Theorem 1 (A & M-G & R)

$$\mathrm{Cyc}(\mathrm{Sp}) \simeq \lim^{r.\mathrm{lax}} \left(\mathrm{Fun}(\mathrm{BT}, \mathrm{Sp}) \underset{\tau}{\overset{\mathrm{lax}}{\curvearrowright}} \mathbb{N}^\times \right).$$

- ★ an object of $\lim^{r.\mathrm{lax}}$ is given by $T \in \mathrm{Fun}(\mathrm{BT}, \mathrm{Sp})$ equipped with:
- for each $r \in \mathbb{N}^\times$, a cyclotomic structure map $T \xrightarrow{\sigma_r} T^{\tau C_r}$;

Theorem 1 (A & M-G & R)

$$\mathrm{Cyc}(\mathrm{Sp}) \simeq \lim^{r.\mathrm{lax}} \left(\mathrm{Fun}(\mathrm{BT}, \mathrm{Sp}) \begin{array}{c} \xrightarrow{\mathrm{lax}} \\ \tau \\ \xleftarrow{\mathrm{lax}} \end{array} \mathbb{N}^\times \right).$$

- ★ an object of $\lim^{r.\mathrm{lax}}$ is given by $T \in \mathrm{Fun}(\mathrm{BT}, \mathrm{Sp})$ equipped with:
- for each $r \in \mathbb{N}^\times$, a cyclotomic structure map $T \xrightarrow{\sigma_r} T^{\tau C_r}$;
 - for each $r, s \in \mathbb{N}^\times$,

Theorem 1 (A & M-G & R)

$$\mathrm{Cyc}(\mathrm{Sp}) \simeq \lim^{r.\mathrm{lax}} \left(\mathrm{Fun}(\mathrm{B}\mathbb{T}, \mathrm{Sp}) \begin{array}{c} \xrightarrow{\mathrm{lax}} \\ \tau \\ \xleftarrow{\mathrm{lax}} \end{array} \mathbb{N}^\times \right).$$

★ an object of $\lim^{r.\mathrm{lax}}$ is given by $T \in \mathrm{Fun}(\mathrm{B}\mathbb{T}, \mathrm{Sp})$ equipped with:

- for each $r \in \mathbb{N}^\times$, a cyclotomic structure map $T \xrightarrow{\sigma_r} T^{\tau C_r}$;
- for each $r, s \in \mathbb{N}^\times$, the *data* of a commutative square

$$\begin{array}{ccc} T & \xrightarrow{\sigma_r} & T^{\tau C_r} \\ \sigma_{rs} \downarrow & & \downarrow (\sigma_s)^{\tau C_r} \\ T^{\tau C_{rs}} & \xrightarrow{\mathrm{can.}} & (T^{\tau C_s})^{\tau C_r} \end{array}$$

Theorem 1 (A & M-G & R)

$$\text{Cyc}(\text{Sp}) \simeq \lim^{r.\text{lax}} \left(\text{Fun}(\text{BT}, \text{Sp}) \underset{\tau}{\overset{\text{l.lax}}{\curvearrowright}} \mathbb{N}^\times \right).$$

★ an object of $\lim^{r.\text{lax}}$ is given by $T \in \text{Fun}(\text{BT}, \text{Sp})$ equipped with:

- for each $r \in \mathbb{N}^\times$, a cyclotomic structure map $T \xrightarrow{\sigma_r} T^{\tau C_r}$;
- for each $r, s \in \mathbb{N}^\times$, the *data* of a commutative square

$$\begin{array}{ccc} T & \xrightarrow{\sigma_r} & T^{\tau C_r} \\ \sigma_{rs} \downarrow & & \downarrow (\sigma_s)^{\tau C_r} \\ T^{\tau C_{rs}} & \xrightarrow{\text{can.}} & (T^{\tau C_s})^{\tau C_r} \end{array}$$

- for each $r_1, \dots, r_n \in \mathbb{N}^\times$,

Theorem 1 (A & M-G & R)

$$\text{Cyc}(\text{Sp}) \simeq \lim^{r.\text{lax}} \left(\text{Fun}(\text{B}\mathbb{T}, \text{Sp}) \underset{\tau}{\overset{\text{l.lax}}{\curvearrowright}} \mathbb{N}^\times \right).$$

★ an object of $\lim^{r.\text{lax}}$ is given by $T \in \text{Fun}(\text{B}\mathbb{T}, \text{Sp})$ equipped with:

- for each $r \in \mathbb{N}^\times$, a cyclotomic structure map $T \xrightarrow{\sigma_r} T^{\tau C_r}$;
- for each $r, s \in \mathbb{N}^\times$, the *data* of a commutative square

$$\begin{array}{ccc} T & \xrightarrow{\sigma_r} & T^{\tau C_r} \\ \sigma_{rs} \downarrow & & \downarrow (\sigma_s)^{\tau C_r} \\ T^{\tau C_{rs}} & \xrightarrow{\text{can.}} & (T^{\tau C_s})^{\tau C_r} \end{array}$$

- for each $r_1, \dots, r_n \in \mathbb{N}^\times$, the *data* of a commutative n -cube...

Theorem 1 (A & M-G & R)

$$\text{Cyc}(\text{Sp}) \simeq \lim^{r.\text{lax}} \left(\text{Fun}(\text{BT}, \text{Sp}) \underset{\tau}{\overset{\text{l.lax}}{\curvearrowright}} \mathbb{N}^\times \right).$$

★ an object of $\lim^{r.\text{lax}}$ is given by $T \in \text{Fun}(\text{BT}, \text{Sp})$ equipped with:

- for each $r \in \mathbb{N}^\times$, a cyclotomic structure map $T \xrightarrow{\sigma_r} T^{\tau C_r}$;
- for each $r, s \in \mathbb{N}^\times$, the *data* of a commutative square

$$\begin{array}{ccc} T & \xrightarrow{\sigma_r} & T^{\tau C_r} \\ \sigma_{rs} \downarrow & & \downarrow (\sigma_s)^{\tau C_r} \\ T^{\tau C_{rs}} & \xrightarrow[\text{can.}]{\sim} & (T^{\tau C_s})^{\tau C_r} \end{array}$$

Thm. [Nikolaus-Scholze]
for T connective and $r=s=p$ prime
Cor.: suff to specify just σ_p
(since $\sigma_{p^n} = (\sigma_p)^{\circ n}$, and n -cubes
canonically commute $\forall n \geq 2$)

- for each $r_1, \dots, r_n \in \mathbb{N}^\times$, the *data* of a commutative n -cube...

Theorem 1 (A & M-G & R)

$$\text{Cyc}(\text{Sp}) \simeq \lim^{r.\text{lax}} \left(\text{Fun}(\text{BT}, \text{Sp}) \underset{\tau}{\overset{\text{l.lax}}{\curvearrowright}} \mathbb{N}^\times \right).$$

★ an object of $\lim^{r.\text{lax}}$ is given by $T \in \text{Fun}(\text{BT}, \text{Sp})$ equipped with:

- for each $r \in \mathbb{N}^\times$, a cyclotomic structure map $T \xrightarrow{\sigma_r} T^{\tau C_r}$;
- for each $r, s \in \mathbb{N}^\times$, the *data* of a commutative square

Thm. [Nikolaus-Scholze]
for T connective and $r=s=p$ prime
Cor.: suff to specify just σ_p
(since $\sigma_{p^n} = (\sigma_p)^{\circ n}$, and n -cubes
canonically commute $\forall n \geq 2$)

$$\begin{array}{ccc} T & \xrightarrow{\sigma_r} & T^{\tau C_r} \\ \sigma_{rs} \downarrow & & \downarrow (\sigma_s)^{\tau C_r} \\ T^{\tau C_{rs}} & \xrightarrow[\text{can.}]{\sim} & (T^{\tau C_s})^{\tau C_r} \end{array}$$

- for each $r_1, \dots, r_n \in \mathbb{N}^\times$, the *data* of a commutative n -cube...

★ **slogan:** $\text{TC}(X)$ is built from $\text{THH}(X) \approx \mathcal{O}(\mathcal{L}X)$ by selecting just those functions:

Theorem 1 (A & M-G & R)

$$\text{Cyc}(\text{Sp}) \simeq \lim^{r.\text{lax}} \left(\text{Fun}(\mathbb{B}\mathbb{T}, \text{Sp}) \underset{\tau}{\overset{\text{l.lax}}{\curvearrowright}} \mathbb{N}^\times \right).$$

★ an object of $\lim^{r.\text{lax}}$ is given by $T \in \text{Fun}(\mathbb{B}\mathbb{T}, \text{Sp})$ equipped with:

- for each $r \in \mathbb{N}^\times$, a cyclotomic structure map $T \xrightarrow{\sigma_r} T^{\tau C_r}$;
- for each $r, s \in \mathbb{N}^\times$, the *data* of a commutative square

Thm. [Nikolaus-Scholze]
for T connective and $r=s=p$ prime
Cor.: suff to specify just σ_p
(since $\sigma_p^n = (\sigma_p)^{\circ n}$, and n -cubes
canonically commute $\forall n \geq 2$)

$$\begin{array}{ccc} T & \xrightarrow{\sigma_r} & T^{\tau C_r} \\ \sigma_{rs} \downarrow & & \downarrow (\sigma_s)^{\tau C_r} \\ T^{\tau C_{rs}} & \xrightarrow[\text{can.}]{\sim} & (T^{\tau C_s})^{\tau C_r} \end{array}$$

- for each $r_1, \dots, r_n \in \mathbb{N}^\times$, the *data* of a commutative n -cube...

★ **slogan:** $\text{TC}(X)$ is built from $\text{THH}(X) \approx \mathcal{O}(\mathcal{L}X)$ by selecting just those functions:

- that are \mathbb{T} -invariant;

Theorem 1 (A & M-G & R)

$$\text{Cyc}(\text{Sp}) \simeq \lim^{r.\text{lax}} \left(\text{Fun}(\mathbb{B}\mathbb{T}, \text{Sp}) \underset{\tau}{\overset{\text{l.lax}}{\curvearrowright}} \mathbb{N}^\times \right).$$

★ an object of $\lim^{r.\text{lax}}$ is given by $T \in \text{Fun}(\mathbb{B}\mathbb{T}, \text{Sp})$ equipped with:

- for each $r \in \mathbb{N}^\times$, a cyclotomic structure map $T \xrightarrow{\sigma_r} T^{\tau C_r}$;
- for each $r, s \in \mathbb{N}^\times$, the *data* of a commutative square

Thm. [Nikolaus–Scholze]
for T connective and $r=s=p$ prime
Cor.: suff to specify just σ_p
(since $\sigma_p^n = (\sigma_p)^{\circ n}$, and n -cubes
canonically commute $\forall n \geq 2$)

$$\begin{array}{ccc} T & \xrightarrow{\sigma_r} & T^{\tau C_r} \\ \sigma_{rs} \downarrow & & \downarrow (\sigma_s)^{\tau C_r} \\ T^{\tau C_{rs}} & \xrightarrow[\text{can.}]{\sim} & (T^{\tau C_s})^{\tau C_r} \end{array}$$

- for each $r_1, \dots, r_n \in \mathbb{N}^\times$, the *data* of a commutative n -cube...

★ **slogan:** $\text{TC}(X)$ is built from $\text{THH}(X) \approx \mathcal{O}(\mathcal{L}X)$ by selecting just those functions:

- that are \mathbb{T} -invariant;
- whose values on $S^1 \xrightarrow{\gamma} X$ determine their values on $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$ “to the greatest extent possible”,

Theorem 1 (A & M-G & R)

$$\text{Cyc}(\text{Sp}) \simeq \lim^{r.\text{lax}} \left(\text{Fun}(\mathbb{B}\mathbb{T}, \text{Sp}) \underset{\tau}{\overset{\text{l.lax}}{\curvearrowright}} \mathbb{N}^\times \right).$$

★ an object of $\lim^{r.\text{lax}}$ is given by $T \in \text{Fun}(\mathbb{B}\mathbb{T}, \text{Sp})$ equipped with:

- for each $r \in \mathbb{N}^\times$, a cyclotomic structure map $T \xrightarrow{\sigma_r} T^{\tau C_r}$;
- for each $r, s \in \mathbb{N}^\times$, the *data* of a commutative square

Thm. [Nikolaus-Scholze]
for T connective and $r=s=p$ prime
Cor.: suff to specify just σ_p
(since $\sigma_p^n = (\sigma_p)^{\circ n}$, and n -cubes
canonically commute $\forall n \geq 2$)

$$\begin{array}{ccc} T & \xrightarrow{\sigma_r} & T^{\tau C_r} \\ \sigma_{rs} \downarrow & & \downarrow (\sigma_s)^{\tau C_r} \\ T^{\tau C_{rs}} & \xrightarrow[\text{can.}]{\sim} & (T^{\tau C_s})^{\tau C_r} \end{array}$$

- for each $r_1, \dots, r_n \in \mathbb{N}^\times$, the *data* of a commutative n -cube...

★ **slogan:** $\text{TC}(X)$ is built from $\text{THH}(X) \approx \mathcal{O}(\mathcal{L}X)$ by selecting just those functions:

- that are \mathbb{T} -invariant;
- whose values on $S^1 \xrightarrow{\gamma} X$ determine their values on $S^1 \xrightarrow{r} S^1 \xrightarrow{\gamma} X$ “to the greatest extent possible”, subject to all possible coherences between these determinations.

Theorem 2 (A & M-G & R)

Theorem 2 (A & M-G & R)

①

diagonal package for spaces

Theorem 2 (A & M-G & R)

①

diagonal package for spaces \rightsquigarrow

$$\begin{array}{ccc} \text{Cat}(\mathcal{S}) & \dashrightarrow & \text{Cyc}^h(\mathcal{S}) \\ & \searrow \text{THH}_{\mathcal{S}} & \downarrow \text{fgt} \\ & & \mathcal{S} \end{array}$$

Theorem 2 (A & M-G & R)

①

diagonal package for spaces \rightsquigarrow

$$\begin{array}{ccc} \text{Cat}(\mathcal{S}) & \dashrightarrow & \text{Cyc}^h(\mathcal{S}) \\ & \searrow \text{THH}_{\mathcal{S}} & \downarrow \text{fgt} \\ & & \mathcal{S} \end{array}$$

$\text{Cyc}^h(\mathcal{S}) :=$ “unstable cyclotomic spaces”

Theorem 2 (A & M-G & R)

①

diagonal package for spaces \rightsquigarrow

$$\begin{array}{ccc} \text{Cat}(\mathcal{S}) & \dashrightarrow & \text{Cyc}^h(\mathcal{S}) \\ & \searrow \text{THH}_{\mathcal{S}} & \downarrow \text{fgt} \\ & & \mathcal{S} \end{array}$$

$\text{Cyc}^h(\mathcal{S}) :=$ “unstable cyclotomic spaces”

②

diagonal package for spaces

linearization
(à la Goodwillie calculus)

$$\rightsquigarrow$$

Tate package for spectra

Theorem 2 (A & M-G & R)

1

diagonal package for spaces \rightsquigarrow

$$\begin{array}{ccc} \text{Cat}(\mathcal{S}) & \dashrightarrow & \text{Cyc}^h(\mathcal{S}) \\ & \searrow \text{THH}_{\mathcal{S}} & \downarrow \text{fgt} \\ & & \mathcal{S} \end{array}$$

$\text{Cyc}^h(\mathcal{S}) :=$ “unstable cyclotomic spaces”

2

diagonal package for spaces

linearization
(à la Goodwillie calculus) \Downarrow
Tate package for spectra \rightsquigarrow

$$\begin{array}{ccc} \text{Cat}(\mathcal{S}p) & \dashrightarrow & \text{Cyc}(\mathcal{S}p) \\ & \searrow \text{THH} & \downarrow \text{fgt} \\ & & \mathcal{S}p \end{array}$$

Theorem 2 (A & M-G & R)

1

diagonal package for spaces \rightsquigarrow

$$\begin{array}{ccc} \text{Cat}(\mathcal{S}) & \dashrightarrow & \text{Cyc}^h(\mathcal{S}) \\ & \searrow \text{THH}_{\mathcal{S}} & \downarrow \text{fgt} \\ & & \mathcal{S} \end{array}$$

$\text{Cyc}^h(\mathcal{S}) :=$ “unstable cyclotomic spaces”

2

diagonal package for spaces

linearization
(à la Goodwillie calculus)

Tate package for spectra \rightsquigarrow

$$\begin{array}{ccc} \text{Cat}(\mathcal{S}p) & \dashrightarrow & \text{Cyc}(\mathcal{S}p) \\ & \searrow \text{THH} & \downarrow \text{fgt} \\ & & \mathcal{S}p \end{array}$$

$\text{Cyc}(\mathcal{S}p) :=$ cyclotomic spectra